



Original Article

 Antifungal activity of extracts and phenolic compounds from *Deguelia duckeana*

 Nerilson M. Lima^{a,1}, Lorena M. Cursino-Hron^{b,1}, Alita M. Lima^b, João V.B. Souza^b,
 André C. de Oliveira^a, Jane V.N. Marinho^a, Cecília V. Nunez^{a,*}
^a Laboratório de Bioprospecção e Biotecnologia, Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brazil

^b Laboratório de Micologia, Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brazil

ARTICLE INFO

Article history:

Received 14 May 2018

Accepted 13 August 2018

Available online 20 September 2018

Keywords:

Amazonian plant

Chalcone

Flavonoid

Lignan

Stilbene

Timbó

ABSTRACT

Candida spp. is associated with almost 80% of all nosocomial fungal infections and is considered a major cause of blood stream infections. In humans, Cryptococcosis is a disease of the lungs caused by the fungi *Cryptococcus gattii* and *Cryptococcus neoformans*. It can be potentially fatal, especially in immune-compromised patients. In a search for antifungal drugs, *Deguelia duckeana* extracts were assayed against these two fungi and also against *Candida albicans*, which causes candidiasis. Hexane branches and CH₂Cl₂ root extracts as well as the substances 4-hydroxy-lonchocarpine, 3,5,4'-trimethoxy-4-prenylstilbene and 3',4'-methylenedioxy-7-methoxyflavone were assayed to determine the minimal inhibitory concentration. Phytochemical study of CH₂Cl₂ root and hexane branch extracts from *D. duckeana* A.M.G. Azevedo, Fabaceae, resulted in the isolation and characterization of nine phenolic compounds: 4-hydroxyderricine, 4-hydroxy-lonchocarpine, 3',4',7-trimethoxy-flavonol, 5,4'-dihydroxy-isolonchocarpine, 4-hydroxyderricine, derricine, 3,5,4'-trimethoxy-stilbene, 3',4',7-trimethoxyflavone and yangambin. The only active extract was a CH₂Cl₂ root showing minimal inhibitory concentration 800 µg/ml against *C. gattii*, and the investigation of compounds obtained from this extract showed that 4-hydroxy-lonchocarpine was active against all three fungi (*C. neoformans*, *C. gattii* and *C. albicans*). These results suggest that *D. duckeana* extracts have potential therapeutic value for the treatment of pathogenic fungi.

© 2018 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Farmacognosia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Candida spp. is associated with almost 80% of all nosocomial fungal infections and is considered the major cause of blood stream infections and its infections involve a broad spectrum of superficial and invasive diseases. The result is a great medical challenge, due to both the difficulties of diagnosis and in finding effective countermeasures to the infections caused by these fungi (Colombo and Guimarães, 2003).

The Fabaceae is a large botanical family and a producer of phenolic compounds such as flavonoids and isoflavonoids used as chemotaxonomic markers (Hegnauer and Gpayer-Barkmeijer, 1993; Veitch, 2013). Species from Papilionoideae subfamily are known to produce substances with pharmacological properties,

including flavonoids from *Tephrosia apollinea* with antifungal activity (Ammar et al., 2013), flavonoids from *Dalbergia odorifera* (Lee et al., 2013), and isoflavonoids from *Abrus mollis*, both with anti-inflammatory activity (Chen et al., 2014a).

Deguelia is one of some 750 genera in the Fabaceae. Studies of the members of this genus (sometimes under synonymies) report stilbene and flavanones from *Derris rariflora* (= *Deguelia rariflora*) (Braz Filho et al., 1975a); rotenone and tephrosin from *Derris urucu* (= *Deguelia rufescens* var. *urucu*) (Braz Filho et al., 1975a); isoflavan from *Derris amazonica* (= *Lonchocarpus negrensis*) (Braz Filho et al., 1975a); stilbene, lonchocarpine and 4-hydroxy-lonchocarpin from *Derris floribunda* (Braz Filho et al., 1975b); stilbene from *Deguelia spruceana* (Garcia et al., 1986); isoflavonoids from *Derris glabrescens* (= *Lonchocarpus densiflorus*) (Monache et al., 1977); prenylated isoflavonoids (Magalhães et al., 2001) and flavanone from *Deguelia hatschbachii* (Magalhães et al., 2003); prenylated flavonoids from *Deguelia longeracemosa* (Magalhães et al., 2006); dihydroflavonols from *D. urucu* (Lôbo et al., 2009), stilbenes from *D. rufescens* var. *urucu* (Lôbo et al., 2010); isoflavonoids and chromones (Lawson

* Corresponding author.

E-mail: cecilia@inpa.gov.br (C.V. Nunez).¹ Both authors contributed equally to the manuscript.

et al., 2008), chalcones and rotenoids from *Lonchocarpus nicou* (Lawson et al., 2010); flavonoids from *Deguelia utilis* (Oliveira et al., 2012) and stilbenes from *D. rufescens* (= *Derris urucu*, *Lonchocarpus urucu*) (Pereira et al., 2012). The main characteristic of this genus and its close relatives is the presence of isoprenyl groups but, as a recent review describes (Marques et al., 2015), it also possesses dimethylchromone and related compounds.

Deguelia duckeana A.M.G. Azevedo, Fabaceae, a species endemic to Brazil, is known as “cipó-cururu” or “timbó” and used by indigenous people to kill fish. It is known only from the Brazilian states of Amazonas, Pará and Rondônia (Camargo and Tozzi, 2017). As far as we know from the available literature, there are only three studies published concerning biological activity and/or chemical isolation of *D. duckeana*. One showing extract antimycobacterial activity (Carrion et al., 2013), another the presence of stilbene and chalcones, *Artemia salina* toxicity and moderate activity against *Staphylococcus aureus* (Lima et al., 2013) and a third describing the isolation of flavones, flavanones, chalcones and stilbene and their effect on cellular viability, AMPK, eEF2, eIF2 and eIF4E (Cursino et al., 2016).

Accordingly, the current study was carried out to enhance knowledge of the chemical and biological potential of *D. duckeana*. First, the antifungal activity of root and branch extracts was evaluated against *Cryptococcus gattii*, *C. neoformans* and *Candida albicans*. Thereafter, phytochemical fractionation of these extracts was performed to obtain pure compounds. As 4-hydroxyonchocarpine is described in the literature with activity against fungi (Mbaveng et al., 2008; Dzoyem et al., 2013; Kuete et al., 2013), this chalcone, together with 3',4'-methylenedioxy-7-methoxyflavone and 3,5,4'-trimethoxy-4-prenylstilbene, all three previously isolated (Cursino et al., 2016), were assayed against *C. albicans* which causes candidiasis, a widespread disease (Chakravarthi and Haleagrahara, 2011), and against *C. gattii* and *C. neoformans* which caused Cryptococcosis, a serious disease—notably in immuno-compromised patients. *C. gattii* also causes meningoencephalitis and other central nervous system and pulmonary-linked diseases, which can often be fatal (Chen et al., 2014b).

Materials and methods

General experimental procedure

Spectral data were obtained from Varian Inova (¹H NMR 500 MHz) and Bruker DRX (¹H NMR 400 MHz). Samples were analyzed using CDCl₃ as solvent and internal standard. Compounds **8** and **9** were also analyzed by LC-MS MicroTOF-QII (Bruker Daltonics), ESI, positive mode and Prominence UFLC (Shimadzu) (DAD) SPD-20A. The SiO₂ 60 chromatography column (230–400 mesh) used was made by Merck, Germany, and the Sephadex LH-20 by Sigma. The solvents MeOH, hexane, EtOAc and CH₂Cl₂ were from Vetec. TLC of SiO₂ (UV254, 0.20 mm, Macherey, Nagel, USA).

Reference fungal strains

Candida albicans (ATCC 36232), *Cryptococcus neoformans* (WM 148, genotype VNI) and *Cryptococcus gattii* (WM 178, genotype VGII) were used as reference material. These strains were kindly supplied by the fungus collection held by Fiocruz-Rio de Janeiro, Brazil, and are now preserved in the microbial collection of the National Institute of Amazon Research (INPA), Manaus, Brazil. The cultures were preserved in mineral oil, and subcultures maintained on Sabouraud medium to ensure purity and viability until testing was performed.

Plant material

Roots and branches of *Deguelia duckeana* A.M.G. Azevedo, Fabaceae, were collected on Praia Dourada (Manaus, Amazonas, Brazil) in September 2005. In order to obtain more plant material to perform the chemical fractionation, a new collection was made in August 2009. Vouchers of both plant materials were deposited in the herbarium of Instituto Federal de Educação do Amazonas (EAFM), as accession numbers 10606 and 10613, respectively.

Plant extraction and substances isolation

Roots were dried and extracted with CH₂Cl₂ as solvent, using an ultrasound bath for 20 min (Unique, São Paulo, Brazil), filtered and the procedure repeated twice. Plant material was then dried and then extracted with methanol (MeOH), and finally with H₂O, with all extractions using the same procedure.

Dichloromethane root extract (8 g) was fractionated in a SiO₂ chromatography column with solvents hexane, CH₂Cl₂, EtOAc and MeOH as gradient. Combined fractions 20–30 obtained as medium polarity (EtOAc) were re-fractionated with CH₂Cl₂, CH₂Cl₂/EtOAc and EtOAc/MeOH. TLC preparative analysis of fraction **5** was eluted with CH₂Cl₂/EtOAc 95:5 and showed compounds in mixture (4.1 mg) as **1** and **2**.

Combined fractions 13–15 were purified by open column chromatography using a Sephadex LH-20 with MeOH as elution system yielding compound **3** and a mixture (128 mg) with compounds **2** (~34%), **4** (~26%) and **5** (~40%). NMR spectral data allowed the correct identification of compounds without isolating them. Relative percentages were calculated in mixture by using ¹H NMR integration signals.

Combined fractions 4–5 (2.8 g) obtained from the first fractionation of CH₂Cl₂ root extract were separated with SiO₂ with the solvents hexane, EtOAc and MeOH yielding 50 fractions. Among them, fraction 39 was analyzed by LC-MS on a C-18 analytic column, using a gradient system with ACN/H₂O (0.1% acetic acid) 20% (0–11 min), 100% (11–12 min), 20% (12–15 min) and flow of 0.4 ml/min. The chromatogram showed two peaks at 5.8 and 6 min, corresponding to *m/z* 313.107121 [M + H]⁺ ion (molecular formula C₁₈H₁₆O₅) for compound **8**, and *m/z* 469.182066 [M + Na]⁺ (molecular formula C₂₄H₃₀O₈) for compound **9**.

In order to identify bioactive flavonoids, the hexanic branch extract (2 g) was fractionated with open column chromatography using SiO₂ with solvents hexane, EtOAc and MeOH as gradient. The combined fractions containing flavonoids was obtained using hexane/EtOAc 9:1 until 1:9 as the elution system, yielding compounds **6** (5 mg) and **7** (3.6 mg).

Fractionation of all samples were monitored by ¹H NMR, UV (254 and 365 nm), with reagents FeCl₃, AlCl₃ and Ce(SO₄)₂.

Antifungal activity


Previously isolated compounds (Cursino et al., 2016) were tested in the current study. Only three compounds (4-hydroxyonchocarpine, 3,5,4'-trimethoxy-4-prenylstilbene and 3',4'-methylenedioxy-7-methoxyflavone) were selected because the first has cytotoxic activity reported in the literature and only they showed enough amount. In addition to these three, hexane branch and CH₂Cl₂ root extracts were also assayed to determine the minimal inhibitory concentration (MIC) as set by the Clinical and Laboratory Standards Institute 2008 (CLSI, 2008). Assays were performed in 96-well plates, each containing 100 μl of each previously diluted substance or extract, plus 100 μl of RPMI 1640 broth medium with substance or extract and 100 μl of diluted microorganism containing 2.5 × 10³ CFU/ml. We evaluated concentrations from 800 to 6.25 μg/ml for plant extracts, and

concentrations from 320 to 0.625 $\mu\text{g/ml}$ for isolated substances. *C. albicans* was incubated at 35 °C for 24 h, and *Cryptococcus gattii* and *C. neoformans* at 35 °C for 72 h. Cultivated fungal strains and RPMI 1640 medium were used as negative controls, and amphotericin B (64 $\mu\text{g/ml}$) as a positive control. Dimethyl sulfoxide was used for compound dilution with final concentration in the bioassay below 1%. MIC values were determined visually after 24 h incubation, as the lowest concentration of drug that resulted in both $\geq 50\%$ inhibition and 100% inhibition of growth relative to the growth of the control, as previously described by the Clinical and Laboratory Standards Institute 2008 (CLSI, 2008).

Results

Compound identifications

Phytochemical study of CH_2Cl_2 root and hexane branch extracts from *D. duckeana* resulted in isolation and characterization of nine phenolic compounds: 4-hydroxyderricine (1), 4-hydroxyonchocarpine (2), 3',4',7-trimethoxy-flavonol (3), 5,4'-dihydroxy-isolonchocarpine (4), 4-hydroxyderricidine (5), derricidine (6), 3,5,4'-trimethoxy-stilbene (7), 3',4',7-trimethoxyflavone (8) and yangambin (9).



Chalcone 4-hydroxyderricine (1): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ : 1.68 (3H, s, Me), 1.80 (3H, s, Me), 3.38 (2H, d, $J=7.2$ Hz, H-1''), 3.91 (3H, s, OMe), 5.23 (1H, m, H-2''), 6.49 (1H, d, $J=9.2$ Hz, H-5'), 6.88 (1H, d, $J=8.8$ Hz, H-3 and H-5), 7.47 (1H, d, $J=15.2$ Hz, H- α), 7.56 (1H, d, $J=8.8$ Hz, H-2 and H-6), 7.79 (1H, d, $J=9.2$ Hz, H-6'), 7.83 (1H, d, $J=15.2$ Hz, H- β), 13.45 (1H, s, 2'-OH). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ : 17.8 (C-5''), 21.6 (C-1''), 25.8 (C-4''), 102.0 (C-5'), 114.7 (C-1'), 115.9 (C-3, C-5), 117.4 (C-3'), 118.3 (C- α), 122.0 (C-2''), 127.7 (C-1), 129.1 (C-6'), 131.9 (C-3''), 130.6 (C-2 and C-6), 144.1 (C- β), 158.0 (C-4), 163.0 (C-2'), 163.2 (C-4'), 192.4 (C=O).

Chalcone 4-hydroxyonchocarpine (2): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 1.45 (6H, s), 5.57 (1H, d, $J=10$ Hz), 6.38 (1H, d, $J=8.5$ Hz, H-5'), 6.74 (1H, d, $J=10$ Hz), 6.89 (2H, d, $J=8.5$ Hz), 7.56 (2H, d, $J=8.5$ Hz), 7.54 (1H, d, $J=15.5$ Hz, H- α), 7.72 (1H, d, $J=8.5$ Hz, H-6'), 7.85 (1H, d, $J=15.5$ Hz, H- β), 13.75 (1H, s).

Flavonol 3',4',7-trimethoxy-flavonol (3): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 3.96 (3H, s, OMe), 3.97 (3H, s, OMe), 3.99 (3H, s, OMe),

6.97 (1H, d, $J=8.0$ Hz), 7.00 (1H, d, $J=2.0$ Hz), 7.02 (1H, dd, $J=9.0$ and 2.0 Hz), 7.39 (1H, d, $J=2.5$), 7.59 (1H, dd, $J=8.5$ and 2.0 Hz), 8.15 (1H, d, $J=8.0$ Hz). $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ : 55.9 (OCH₃), 56.0 (OCH₃), 56.3 (OCH₃), 100.1 (C-8), 110.7 (C-5'), 112.2 (C-2'), 115.0 (C-6), 116.2 (C-10), 123.1 (C-6'), 124.0 (C-1'), 127.8 (C-5), 148.6 (C-3'), 151.3 (C-4'), 152.4 (C-9), 160.0 (C-2), 164.5 (C-7), 172.5 (C=O).

5,4'-Dihydroxy-isolonchocarpine (4): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 12.27 (1H, s), 7.30 (2H, dd, $J=8.5$ and 2.5 Hz; H-2' and H-6'), 6.88 (2H, dd, $J=8.5$ and 2.5 Hz; H-3' and H-5'), 6.61 (1H, d, $J=10.0$, H-4''), 5.49 (1H, d, $J=10.0$, H-3'') and 1.43 (6H, s), 5.95 (1H, s), 5.32 (1H, dd, $J=13$ and 3.0 Hz, H-2), 3.06 (1H, dd, $J=17.0$ and 13.0 Hz, H-3), 2.77 (1H, dd, $J=17.0$ and 3.0 Hz, H-3).

Chalcone 4-hydroxyderricidine (5): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 6.88 (2H, d, $J=8.5$, H-3 and H-5), 7.54 (2H, d, $J=8.5$, H-2 and H-6), 7.42 (1H, d, $J=15.5$, H- α), 7.83 (1H, d, $J=15.5$, H- β), 7.81 (1H, d, $J=8.5$, H-6'), 6.47 (1H, d, $J=2.5$, H-3'), 6.49 (1H, dd, $J=8.5$ and 2.5, H-5'), 4.56 (2H, d, $J=7.0$, H-1''), 5.49 (1H, m, H-2''), 1.80 (3H, s, CH₃), 1.75 (3H, s, CH₃), 13.56 (1H, s).

Chalcone derricidine (6): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 13.44 (1H, s), 7.89 (1H, d, $J=16.0$ Hz), 7.59 (1H, d, $J=16.0$ Hz), 7.65 (2H, m, H-2 and H-6), 7.44 (3H, m, H-3, H-4 and H-6), 1.81 (3H, s), 1.76 (3H, s),

4.57 (2H, d, $J=6.8$ Hz), 5.49 (1H, m), 6.50 (1H, dd, $J=8.4$ and 2.4 Hz), 6.48 (1H, d, $J=2.4$ Hz), 7.83 (1H, d, $J=8.4$ Hz).

Stilbene 3,5,4'-trimethoxy-stilbene (7): $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 7.04 (1H, d, $J=16.4$ Hz, H-8), 6.91 (1H, d, $J=16.4$ Hz, H-7), 3.83 (3H, s), 7.45 (2H, dd, $J=8.4$ and 2.0 Hz, H-2' and H-6'), 6.90 (2H, d, $J=8.4$ and 2.0 Hz, H-3' and H-5'), 3.832 (6H, s), 6.65 (2H, d, $J=2.0$ Hz, H-2 and H-6), 6.37 (1H, t, $J=2.0$ Hz, H-4).

Flavone 3',4',7-trimethoxyflavone (8): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ : 3.92 (3H, s, 7-OCH₃), 3.95 (3H, s, 4'-OCH₃), 3.97 (3H, s, 3'-OCH₃), 6.98 (1H, d, $J=2.0$, H-8), 6.98 (2H, dd, $J=8.5$ and 2.0, H-6), 8.11 (1H, d, $J=8.5$, H-5), 6.69 (1H, s, H-3), 7.53 (1H, dd, $J=8.5$ and 2.0, H-6'), 6.96 (1H, d, $J=8.5$, H-5'), 7.35 (1H, d, $J=2.0$, H-2'). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ : 55.8 (7-OCH₃), 56.0 (4'-OCH₃), 56.1 (3'-OCH₃), 117.6 (C-10), 157.8 (C-9), 111.0 (C-8), 164.0 (C-7), 114.2 (C-6), 126.9 (C-5), 177.8 (C-4), 106.3 (C-3), 163.0 (C-2), 119.8 (C-6'), 100.3 (C-5'), 151.8 (C-4'), 149.2 (C-3'), 108.7 (C-2'), 124.2 (C-1').

Table 1
MIC ($\mu\text{g/ml}$) of extracts and isolated compounds from *Deguelia duckeana* against *Candida albicans*, *Cryptococcus gattii* and *Cryptococcus neoformans*.

	<i>C. albicans</i> (ATCC 36232)	<i>C. gattii</i> (WM 17)	<i>C. neoformans</i> (WM 148)
CH ₂ Cl ₂ root extract	>800	800	>800
Hexanic branch extract	>800	>800	>800
4-Hydroxy lonchocarpine	320	80	20
3',4'-Methylenedioxy-7-methoxyflavone	>320	>320	>320
3,5,4'-Trimethoxy-4-prenylstilbene	>320	>320	>320
Amphotericin B	0.25	0.125	0.06

Lignan yangambin (**9**): ¹H NMR (400 MHz, CDCl₃) δ : 3.09 (2H, m, 8, 8'), 3.82 (6H, s, OCH₃), 3.86 (12H, s, OCH₃), 3.92 (2H, dd, $J=9.0$ and 6.9 Hz, H-9 β , H-9' β), 4.31 (2H, dd, $J=9.0$ and 6.9 Hz, H-9 α , H-9' α), 4.75 (2H, d, $J=4.0$, H-7, 7'), 6.57 (4H, s, H-2, H-6, H-2', H-6'), ¹³C NMR (100 MHz, CDCl₃) δ : 54.3 (C-8, C-8'), 55.8 (OCH₃), 60.8 (OCH₃), 71.9 (C-9, C-9'), 85.9 (C-7, C-7'), 102.7 (C-2, C-6, C-2', C-6'), 136.6 (C-1, C-1'), 137.4 (C-4, C-4'), 153.3 (C-3, C-3', C-5, C-5').

Although all these compounds are known, it is important to emphasize that *D. duckeana* has been reported as a species with important biological activities, but is so far very under-researched. The current study of *D. duckeana* found several phenolic compounds which corroborate known Fabaceae chemotaxonomy.

Antifungal activity

In terms of the MIC, as set by the Clinical and Laboratory Standards Institute 2008 (CLSI, 2008), the antifungal activity of 4-hydroxy lonchocarpine showed significant results for members of the genera *Candida* and *Microsporum*, but the activity against *C. gattii*, described in the current study (Table 1), is being done so for first time, as far as we know.

Discussion

The identification of nine phenolic compounds from *D. duckeana* by the current study contributes to chemosystematic knowledge of genus *Deguelia*, which shows mostly flavonoid and related compounds. The compounds 4-hydroxy lonchocarpine and derricidine were previously isolated from *D. duckeana* branches (Braz Filho et al., 1975b; Oliveira et al., 2012; Lima et al., 2013; Cursino et al., 2016; Ahmed et al., 2002), while 4-hydroxyderricidine is described for the first time from the genus *Deguelia*.

Compound **1** (4-hydroxyderricidine) was a yellow solid in mixture with compound **2** (4-hydroxy lonchocarpine). Integration analysis of the signals from spectral data allowed the identification of compound **2** corresponding to 47.7% of the mixture. Two singlets at δ_{H} 13.45 and δ_{H} 13.75 indicate the presence of two flavonoids with chelated hydroxyl groups. Compound **1** was characterized as a chalcone due to two doublets at δ_{H} 7.47 (1H, $J=15.2$ Hz) and δ_{H} 7.83 (1H, $J=15.2$ Hz) related to the trans-olefinic hydrogens α and β , one singlet in δ_{H} 13.45 referable to a chelated hydroxyl group (C2'-OH), two doublets at δ_{H} 7.56 (2H, $J=8.8$ Hz) and 6.88 (2H, $J=8.8$ Hz) corresponding to H-2/H-6 and H-3/H-5, respectively. Two doublets at δ_{H} 7.79 (1H, $J=9.2$ Hz, H-6') and δ_{H} 6.49 (1H, $J=9.2$ Hz, H-5') indicated the presence of *ortho* coupling and one singlet at δ_{H} 3.91 (3H) characterized a methoxyl linked to an aromatic group. At δ_{H} 5.23 (1H, m), δ_{H} 3.38 (2H, d, $J=7.2$ Hz), 1.80 (3H, s) and 1.68 (3H, s), a prenyl group was observed. The chemical shifts of carbon 4 (δ_{C} 158.0), 3 and 5 (δ_{C} 115.9) indicated that there was only a hydroxyl group linked to carbon 4. The structural proposal was confirmed by comparison with the literature (Shin et al., 2011).

Compound **2**, identified as chalcone 4-dihydroxy lonchocarpine, is common in Fabaceae and Moraceae families. Chalcones are known to possess antimalarial (Ramírez et al., 2010), antibacterial,

antifungal (Dzoyem et al., 2013) and anticancer (Ngameni et al., 2006) biological activity.

¹H NMR spectrum of compound **3** (3',4',7-trimethoxy-flavonol) showed *ortho* and *meta* hydrogens coupled with double doublets at δ_{H} 7.03 ($J=8.9$ and 2.4 Hz) and 6.91 ($J=2.4$ Hz) and a doublet at δ_{H} 8.20 (1H, $J=8.9$ Hz) characterizing H-6, H-5 and H-8 of A-ring belonging to flavonoid nuclei, respectively.

In the ¹³C NMR spectrum, 15 carbon sp² were observed, compatible with units of the C6-C3-C6 typical of flavonoids. The signal at δ_{C} 172.5 (C-4) is compatible with a flavonoid carbonyl group. The signals δ_{C} 112.2, 123.1 and 110.7 correlated on an HSQC contour map with δ_{H} 7.45 (d, $J=2.1$ Hz), 7.57 (dd, $J=8.4$ and 2.1 Hz) and 7.01 (d, $J=8.4$ Hz), respectively, and indicated a B ring at the C-3 and C-4 positions. Verified signals δ_{C} 55.9, 56.0 and 56.3 on ¹³C NMR correlated with singlets at δ_{H} 3.93, 3.97 and 3.98 on ¹H NMR and indicated the presence of three methoxyl aromatic groups which were assigned to C-7, C-3 and C-4 carbons. Localization of methoxyl substituent groups was confirmed by an HMBC contour map.

Compound **4** (5,4'-dihydroxy-isolonchocarpin) was recognized as a flavanone through signals of C-ring hydrogens [δ_{H} 5.32 (1H, dd, $J=13.0$ and 3.0 Hz, H-2), 3.06 (1H, dd, $J=17.0$ and 13.0 Hz, H-3), 2.77 (1H, dd, $J=17.0$ and 3.0 Hz, H-3)]. One singlet at δ_{C} 12.27 characterized a chelated hydroxyl group that could be assigned to a flavanone C-5. Signals from a *para*-substituted B-ring [δ_{H} 7.30 (2H, dd, $J=8.5$ and 2.5 Hz, H-2' and H-6') and 6.88 (2H, dd, $J=8.5$ and 2.5 Hz, H-3' and H-5')], signals of gem-dimethyl-chromone [δ_{H} 6.61 (1H, d, $J=10.0$, H-4''), 5.49 (1H, d, $J=10.0$, H-3'') and 1.43 (6H, s)], and one singlet at H-8 [δ_{H} 5.95 (1H; s)] characterized the flavanone 5,4'-dihydroxy-isolonchocarpin.

Compound **5** (4-hydroxyderricidine) showed a doublet at 6.88 (2H, $J=8.5$, H-3 and H-5) and 7.54 (2H, d, $J=8.5$, H-2 and H-6), indicating B ring substitution by a chelated hydroxyl group at δ_{H} 13.56.

Compound **6** (derricidine) was obtained as a yellow solid, a precursor of 4-hydroxyderricidine (**5**). ¹H NMR spectrum of compound **6** revealed the presence of two double doublets at δ_{H} 7.89 and 7.59 ($J=16.0$ Hz), characterizing the *trans*-olefinic system of chalcone. A singlet at δ_{H} 13.44 indicated a chelated hydroxyl group on the C-2' position. The signals at δ_{H} 7.65 (2H, m, H-2 and H-6) and 7.44 (3H, m, H-3, H-4 and H-6) indicated an unsubstituted B ring. A prenyl group was observed via the signals at 1.81 (s, 3H), 1.76 (s, 3H), 4.57 (2H, d, $J=6.8$ Hz) and 5.49, and also at 6.50 (1H, $J=8.4$ and 2.4 Hz) with *meta* coupling at 6.48 (1H, $J=2.4$ Hz) and *ortho* coupling at 7.83 (1H, d, $J=8.4$ Hz).

Compound **7** was characterized as a trimethoxylated derivative of resveratrol (3,5,4'-trimethoxy-stilbene). It showed doublets at δ_{H} 7.04 and 6.91 with large J -coupling (16.4 Hz) characterizing *trans*-ethylenic chair due to two singlets of aromatic methoxyl at δ_{H} 3.833 (3H) and δ_{H} 3.832 (6H), three singlets of aromatic methoxyl and two pairs of doublets δ_{H} 7.45 (2H, H-2' and H-6') and δ_{H} 6.90 (2H, H-3' and H-5') with coupling at *ortho* ($J=8.4$ Hz) and *meta* ($J=2.0$ Hz), indicating one *para*-substituted aromatic system. The methoxyl groups were attributed to the 3 and 5 positions owing to two doublets with coupling *meta* at δ_{H} 6.65 (2H, $J=2.0$ Hz) and one triplet at δ_{H} 6.37 attributed to homotopic hydrogens H-2, H-6 and to H-4, respectively.

Compounds **8** and **9** were identified in mixture. ^1H NMR spectrum of compound **8** (3',4',7-trimethoxyflavone) showed one singlet at δ_{H} 6.69, indicating the presence of hydrogen of a flavone C-ring. The signals at δ_{H} 8.11 (1H, *d*, *J* = 8.5 Hz, H-5), 6.98 (1H, *dd*, *J* = 8.5 and 2.0 Hz, H-6) and 6.98 (1H, *d*, *J* = 2.0 Hz, H-6) belong to the hydrogens H-5, H-6 and H-8 (A-ring). Chemical shifts from a B ring were observed at δ_{H} 7.53 (*dd*, *J* = 8.5 and 2.0, H-6'), 6.96 (*d*, *J* = 8.5, H-5'), 7.35 (*d*, *J* = 2.0, H-2'), indicating substitutions on C-3' and C-4' positions due to methoxyl groups at δ_{H} 3.95 (*s*, 3H) and 3.97 (*s*, 3H). A methoxyl group was observed at δ_{H} 3.92 (*s*) attributed to C-7, which was confirmed through correlations on a contour map. Combined, these spectral data allowed the identification of 3',4',7-trimethoxyflavone.

Compound **9** was identified as the lignan yangambin through signals at δ_{H} 6.57 (4H, *s*), δ_{H} 4.75 (2H, *d*, *J* = 4.0 Hz), 4.31 (2H, *dd*, *J* = 9.0 and 6.9 Hz), 3.92 (2H, *dd*, *J* = 9.0 and 6.9) and δ 3.09 (2H, *m*). These signals showed correlation on an HSQC contour map: δ_{C} 102.7 and δ_{H} 6.57; δ_{C} 85.9 and δ_{H} 4.75; δ_{C} 54.3 and δ_{H} 3.09 and on HMBC: δ_{C} 137.4 and δ_{H} 6.57 (3J); δ_{C} 137.4 and δ_{H} 3.82 (3J); δ_{C} 85.9 and δ_{H} 6.57 (3J); δ_{C} 85.9 and δ_{H} 4.31 (3J); δ_{C} 54.3 and δ_{H} 4.75 (2J).

In order to determine minimum inhibition concentrations (MICs), hexanic branches and CH_2Cl_2 root extracts were tested against *C. albicans*, *C. gattii* and *C. neoformans*. The only active extract was CH_2Cl_2 root, which showed an MIC of 800 $\mu\text{g}/\text{ml}$ against *C. gattii*. Investigation of the compounds obtained from this extract showed that 4-hydroxyronchocarpine was active against all three species. 4-Hydroxyronchocarpine has been described as a chalcone with several biological activities, including antimicrobial, anticancer, antituberculosis, antimalarial, antioxidant and anti-inflammatory potential (Kuetze et al., 2013) and, more recently, induced lactate dehydrogenase (LDH), phosphorylation of the eukaryotic elongation factor 2 (eEF2) and AMP-activated protein kinase (AMPK) and activated caspase-3 (Cursino et al., 2016).

The antifungal activity of 4-hydroxyronchocarpine showed significant results for *C. albicans*, *C. gabrata*, *Microsporum audouinii* and *Trichophyton rubrum* (Mbaveng et al., 2008), and against *C. tropicalis*, *C. albicans* and *C. neoformans* (Dzoyem et al., 2013), but that for *C. gattii*, described in the current study (Table 1), is being done so for first time, as far as we know.

The properties of compound **8** (3',4',7-trimethoxyflavone) have already been investigated by another study, which isolated it and tested its effects on phosphorylation of eEF2, AMPK and eIF4E (Cursino et al., 2016). Compound **9** (yangambin) has been described from a wide variety of species, including *Achillea holosericea*, Asteraceae (Ahmed et al., 2002), *Magnolia fargesii*, Magnoliaceae (Kim et al., 2009), *Ocotea duckei*, Lauraceae (Antunes et al., 2006), as well as previous studies of *D. duckeana* (Cursino et al., 2016). It showed analgesic and anticancer activities (Hausott et al., 2003) and protective effect related to cardiovascular collapse (Araújo et al., 2001).

As this genus is known for the presence of prenylated flavonoids, the present results corroborate the location of the *Deguelia* within the Fabaceae. The present study describes antifungal activity of 4-hydroxyronchocarpine against *C. gattii* for the first time, which indicates a preliminary antifungal activity. Further studies, especially in the pharmacological area, are necessary to confirm these results.

Authors' contributions

NML, LMCC, ACO, JVNM, CVN and JVBS conceived and designed the experiments; NML and JVNM collected the plant sample and made herbarium exsiccates; NML, LMCC, AML, ACO and JVNM performed the experiments; NML, LMCC, AML, JVBS and CVN analyzed the data; JVBS and CVN supervised the laboratory work and contributed with reagents/materials/analysis tools, as well as to critical

reading of the manuscript; NML, LMCC and CVN wrote the paper. All the authors have read the final manuscript and approved the submission.

Conflicts of interest

All authors have none to declare.

Acknowledgments

The authors would like to thank to Brazilian Research Agencies CNPq (PPBio/CNPq - 457472/2012-0, CT-Agro/CNPq - 405804/2013-0, REPENSA/CNPq/FAPEAM - 562892/2010-9) and CAPES (Pro-Amazônia/CAPES - 23038.000738/2013-78) for financial support. The authors also thank Andersson Barison and Kahlil Salomé, from University of Paraná, the INPA's Natural Products Analytical Central for the NMR spectra analysis, and Adrian Barnett who helped with the English.

References

- Ahmed, A.A., Mahmoud, A.A., Ali, E.T., Tzakou, O., Couladis, M., Mabry, T.J., Gáti, T., Tóth, G., 2002. Two highly oxygenated eudesmanes and ten lignans from *Achillea holosericea*. *Phytochemistry* 59, 851–856.
- Ammar, M.I., Nenaah, G.E., Mohamed, A.H., 2013. Antifungal activity of prenylated flavonoids isolated from *Tephrosia apollinea* L. against four phytopathogenic fungi. *Crop. Prot.* 49, 21–25.
- Antunes, R.M.P., Lima, E.O., Pereira, M.S.V., Camara, C.A., Arruda, T.A., Catão, R.M.R., Barbosa, T.P., Nunes, X.P., Silva, T.M.S., 2006. Atividade antimicrobiana *in vitro* e determinação da concentração inibitória mínima (CIM) de fitoconstituintes e produtos sintéticos sobre bactérias e fungos leveduriformes. *Rev. Bras. Farmacogn.* 16, 517–524.
- Araújo, C.V., Barbosa-Filho, J.M., Cordeiro, R.S.B., Tibiriçá, E., 2001. Protective effects of yangambin on cardiovascular hyporeactivity to catecholamines in rats with endotoxin-induced shock. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 363, 267–275.
- Braz Filho, R., Gottlieb, O.R., Mourão, A.P., 1975a. A stilbene and two flavanones from *Derris rariflora*. *Phytochemistry* 14, 261–263.
- Braz Filho, R., Gottlieb, O.R., Mourão, A.P., Da Rocha, A.I., Oliveira, F.S., 1975b. Flavonoids from *Derris* species. *Phytochemistry* 14, 1454–1456.
- Camargo, R.A., Tozzi, A.M.G.A., 2017. *Deguelia*, in Flora do Brasil 2020 em construção. Jardim Botânico do Rio de Janeiro. Available from: <http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB83028> (accessed 6 May 2018).
- Carrion, L.L., Ramos, D.F., Martins, D., Osório, M.I.C., Cursino, L.M.C., Mesquita, D.W.O., Nunez, C.V., Silva, P.E.A., 2013. Antimycobacterial activity of Brazilian Amazon plants extracts. *Int. J. Phytomed.* 5, 479–485. <http://dx.doi.org/10.5138/ijpm.v5i4.1202>.
- Chakravarthi, S., Haleagrahara, N., 2011. A comprehensive review of the occurrence and management of systemic candidiasis as an opportunistic infection. *Microbiol. J.* 1, 1–7.
- Chen, M., Wang, T., Jiang, Z., Shan, C., Wang, H., Wu, M., Zhang, S., Zhang, Y., Zhang, L., 2014a. Anti-inflammatory and hepatoprotective effects of total flavonoid C-glycosides from *Abrus mollis* extracts. *Chin. J. Nat. Med.* 12, 590–598.
- Chen, S.C.A., Meyer, W., Sorrell, T.C., 2014b. *Cryptococcus gattii* Infections. *Clin. Microbiol. Rev.* 27, 980–1024.
- CLSI, 2008. Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard, 3rd ed., Clinical and Laboratory Standards Institute – CLSI document M27-A3, Wayne, PA.
- Colombo, A.L., Guimarães, T., 2003. Epidemiology of hematogenous infections due to *Candida* spp. *Rev. Soc. Bras. Med. Trop.* 36, 599–607.
- Cursino, L.M.C., Lima, N.M., Murillo, R., Nunez, C.V., Merfort, I., Humar, M., 2016. Isolation of flavonoids from *Deguelia duckeana* and their effect on cellular viability, AMPK, eEF2, eIF2 and eIF4E. *Molecules* 21, 192–203.
- Dzoyem, J.P., Hamamoto, H., Ngameni, B., Ngadjui, B.T., Sekimizu, K., 2013. Antimicrobial action mechanism of flavonoids from *Dorstenia* species. *Drug Discov. Ther.* 7, 66–72.
- Garcia, M., Kand, M.H.C., Vieira, D.M., Do Nascimento, M.C., Mors, W.B., 1986. Isoflavonoids from *Derris spruceana*. *Phytochemistry* 25, 2425–2427.
- Hausott, B., Greger, H., Marian, B., 2003. Naturally occurring lignans efficiently induce apoptosis in colorectal tumor cells. *J. Cancer Res. Clin. Oncol.* 129, 569–576.
- Hegnauer, R., Gpayer-Barkmeijer, R.J., 1993. Relevance of seed polysaccharides and flavonoids for the classification of the Leguminosae: a chemotaxonomic approach. *Phytochemistry* 4, 3–16.
- Kim, J.Y., Lim, H.J., Lee, D.Y., Kim, J.S., Kim, D.H., Kee, H.J., Kim, H.D., Jeon, R., Ryu, J., 2009. *In vitro* anti-inflammatory activity of lignans isolated from *Magnolia fargesii*. *Bioorg. Med. Chem. Lett.* 19, 937–940.

- Kuete, V., Noumedem, J.A.K., Nana, F., 2013. Chemistry and pharmacology of 4-hydroxyonchocarpin: a review. *Chin. J. Integr. Med.* 19, 475–480.
- Lawson, M.A., Kaouadji, M., Chulia, A.J., 2008. Nor-dehydrodeguelin and nor-dehydrorotenone, C₂₂ coumaronochromones from *Lonchocarpus nicou*. *Tetrahedron Lett.* 49, 2407–2409.
- Lawson, M.A., Kaouadji, M., Chulia, A.J., 2010. A single chalcone and additional rotenoids from *Lonchocarpus nicou*. *Tetrahedron Lett.* 51, 6116–6119.
- Lee, D., Li, B., Keo, S., Kim, K., Jeong, G., Oh, H., Kim, Y., 2013. Inhibitory effect of 9-hydroxy-6,7-dimethoxydalbergiquinol from *Dalbergia odorifera* on the NF- κ B-related neuroinflammatory response in lipopolysaccharide-stimulated mouse BV2 microglial cells is mediated by heme oxygenase-1. *Int. Immunopharmacol.* 17, 828–835.
- Lima, N.M., Andrade, J.I.A., Lima, K.C.S., Santos, F.N., Barison, A., Salomé, K.S., Matsuura, T., Nunez, C.V., 2013. Chemical profile and biological activities of *Deguelia duckeana* A.M.G. Azevedo (Fabaceae). *Nat. Prod. Res.* 27, 425–432.
- Lôbo, L.T., Silva, G.A., Ferreira, M., Silva, M.N., Santos, A.S., Arruda, A.C., Guilhon, G.M.S.P., Santos, L.S., Borges, R.S., Arruda, M.S.P., 2009. Dihydroflavonols from the leaves of *Derris urucu* (Leguminosae): structural elucidation and DPPH radical-scavenging activity. *J. Braz. Chem. Soc.* 20, 1082–1088.
- Lôbo, L.T., Silva, G.A., Freitas, M.C.C., Souza Filho, A.P.S., Silva, M.N., Arruda, A.C., Guilhon, G.M.S.P., Santos, L.S., Santos, A.S., Arruda, M.S.P., 2010. Stilbenes from *Deguelia rufescens* var. *urucu* (Ducke) A.M.G. Azevedo leaves: Effects on seed germination and plant growth. *J. Braz. Chem. Soc.* 21, 1838–1844.
- Magalhães, A.F., Tozzi, A.M.G.A., Magalhães, E.G., Moraes, V.R.S., 2001. Prenylated flavonoids from *Deguelia hatschbachii* and their systematic significance in *Deguelia*. *Phytochemistry* 57, 77–89.
- Magalhães, A.F., Tozzi, A.M.G.A., Magalhães, E.G., Moraes, V.R.S., 2003. New spectral data of some flavonoids from *Deguelia hatschbachii* A.M.G. Azevedo. *J. Braz. Chem. Soc.* 14, 133–137.
- Magalhães, A.F., Tozzi, A.M.G.A., Magalhães, E.G., Souza-Neta, L.C., 2006. New prenylated metabolites of *Deguelia longeracemosa* and evaluation of their antimicrobial potential. *Planta Med.* 72, 358–363.
- Marques, E.J., Serafim, J.C.R.B., Lemes, B.B., Carvalho, M.F.A., Pereira, M.G., Souza Neta, L.C., 2015. Occurrence and distribution of polyphenolics in species of *Deguelia* (Leguminosae). *J. Microb. Biochem. Technol.* 7, 327–333.
- Mbaveng, A.T., Ngameni, B., Kuete, V., Simo, I.K., Ambassa, P., Roy, R., Bezabih, M., Etoa, F., Ngadjui, B.T., Abegaz, B.M., Meyer, J.J.M., Lall, N., Beng, V.P., 2008. Antimicrobial activity of the crude extracts and five flavonoids from the twigs of *Dorstenia barteri* (Moraceae). *J. Ethnopharmacol.* 116, 483–489.
- Monache, F.D., Valera, G.C., Zapata, D.S., Marini-Bettólo, G.B., 1977. 3-Aryl-4-methoxycoumarins and isoflavones from *Derris glabrescens*. *Gazz. Chim. Ital.* 107, 403–407.
- Ngameni, B., Touaibia, M., Patnam, R., Belkaid, A., Sonna, P., Ngadjui, B.T., Annabi, B., Roy, R., 2006. Inhibition of MMP-2 secretion from brain tumor cells suggests chemopreventive properties of a furanocoumarin glycoside and of chalcones isolated from the twigs of *Dorstenia turbinata*. *Phytochemistry* 67, 2573–2579.
- Oliveira, D.G., Almeida, C.M.C., Silva, C.Y.Y., Arruda, M.S.P., Arruda, A.C., Lopes, D.C.F., Yamada, E.S., Costa, E.T., Silva, M.N., 2012. Flavonoids from the leaves of *Deguelia utilis* (Leguminosae): structural elucidation and neuroprotective properties. *J. Braz. Chem. Soc.* 23, 1933–1939.
- Pereira, A.C., Arruda, M.S.P., Silva, E.A.S., Silva, M.N., Lemos, V.S., Cortes, S.F., 2012. Inhibition of α -glucosidase and hypoglycemic effect of stilbenes from the Amazonian plant *Deguelia rufescens* var. *urucu* (Ducke) A.M.G. Azevedo (Leguminosae). *Planta Med.* 78, 36–38.
- Ramírez, M.E., Mendoza, A.J.A., Arreola, G.R.H., Ordaz, P.C., 2010. Flavonoides con actividad antiprotozoaria. *Rev. Mexicana Cien. Farm.* 41, 6–21.
- Shin, J.E., Choi, E.J., Jin, Q., Jin, H., Woo, E., 2011. Chalcones isolated from *Angelica keiskei* and their inhibition of IL-6 production in TNF- α -stimulated MG-63 cell. *Arch. Pharm. Res.* 34, 437–442.
- Veitch, N.C., 2013. Isoflavonoids of the Leguminosae. *Nat. Prod. Rep.* 30, 988–1027.