



Adaptations of Pre-Columbian Manioc Storage Techniques as Strategies to Adapt to Extreme Climatic Events in Amazonian Floodplains

Julia Vieira da Cunha Ávila^{1,2} · Anderson Márcio Amaral^{2,8} · Angela May Steward^{2,3} · André Braga Junqueira⁴ · Gilton Mendes dos Santos⁵ · Tamara Ticktin⁶ · Charles R. Clement⁷

Accepted: 2 September 2022 / Published online: 27 September 2022

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

Abstract

In Amazonian floodplains, manioc flour is the main plant food product and source of income for local populations. In the context of climate change, extreme flooding is more frequent and intense, making it difficult to cultivate and process manioc. As local knowledge is dynamic and fundamental to adapt in critical times, we studied local techniques for storing manioc roots, which allow them to be processed later. We conducted semi-structured interviews in three floodplains (*várzea*) communities (36 families) and three paleo-floodplain (*paleovárzea*; 1–3 m higher) communities (52 families) in the middle Solimões River basin (Brazil). Residents mention four techniques for storing fresh manioc; two were cited in archaeological or ethnographic studies (burial and basketing), and two were not cited before in the region (bagging and *kanaká*). In the paleovárzea, where manioc production is more important as a source of income, residents have more knowledge of manioc storage techniques. However, this knowledge also persists in areas where manioc has less importance for income generation. Residents of the study area express demand for disseminating these practices, as they can contribute to adaptation in critical periods.

Keywords *Ribeirinhos* · Traditional Ecological Knowledge · adaptive strategies · Indian bread · Brazil

Introduction

In Amazonia, local communities base their diet on fresh foods, from fishing, hunting, and harvesting wild or cultivated fruits and vegetables (Alencar, 2010; Dufour et al., 2016; Posey, 1985). In addition, local communities use traditional food processing, storage, and preservation techniques, such as the production of manioc flour (*Manihot esculenta* Crantz) (Denevan, 1996; Lentz, 2000), the leading plant food of current Amazonian populations and an important food source for past populations (Dufour, 1995; Dufour et al., 2016). In climate change scenarios, where extreme floods have become more frequent and intense (Cai et al., 2014; Marengo et al., 2013), local communities' livelihoods and diets have suffered changes (Ávila et al., 2021; Dubreuil et al., 2017; Funatsu et al., 2019; Langill & Abizaid, 2020; Tregidgo et al., 2020). In this study, we seek to identify the strategies used for manioc storage in the context of extreme floods in Amazonian *várzea* and *paleovárzea* ecosystems.

✉ Julia Vieira da Cunha Ávila
biojuba@gmail.com

¹ Graduate Program in Botany, National Research Institute for Amazonia, Avenida André Araújo, 2936, 69067-375 Petrópolis, Manaus, Amazonas, Brazil

² Mamirauá Sustainable Development Institute, Tefé, Amazonas, Brazil

³ Federal University of Pará, Belém, Pará, Brazil

⁴ Autonomous University of Barcelona, Bellaterra, Barcelona, Spain

⁵ Federal University of Amazonas, Manaus, Amazonas, Brazil

⁶ University of Hawaii, Honolulu, Hawaii, USA

⁷ National Research Institute for Amazonia, Manaus, Amazonas, Brazil

⁸ Graduate Studies of Sociocultural Diversity, Emílio Goeldi Museum, Belém, Pará, Brazil

Dry manioc flour is a primary source of carbohydrates (Dufour et al., 2016) in the Amazonian *várzeas* and *paleovárzeas* of the middle Solimões River basin. Dry flour is also local populations' main form of manioc storage (Lima et al., 2012). The region has already documented diverse methods for processing and consuming manioc (Venturato & Pereira, 2010). After harvest, the manioc flour produced is often the basis of family food security throughout the year; additionally, surpluses are sold locally (Adams et al., 2009; Fraser et al., 2012). The durability of manioc flour is also advantageous because in much of Amazonia refrigerating food is not viable (Tregidgo et al., 2020). Limitations include the high electricity costs, coming mainly from generators powered by fossil fuels and, rarely, with solar energy (Penteado et al., 2019; Valer et al., 2014), and refrigerators and freezers.

In Amazonia, the *várzeas* are flooded annually by white water rivers. At the same time, the *paleovárzeas* are only inundated in extreme flood years (Irion et al., 2010) because they are 1–3 m above the *várzeas*. Hence, the impacts of large floods between the *várzea* and *paleovárzea* agroecosystems are different and more intense in the latter environment (Ávila et al., 2021). Because manioc plants cannot withstand flooding for many days (Langill & Abizaid, 2020), local producers quickly seek to process the roots in local flour mills. However, these processing areas can also be flooded. Even if manioc flour mills are mounted on floating rafts, the community demand may be too high for everyone to process (Ávila et al., 2021).

Archaeological and ethnographic studies carried out in *terra firme* (non-flooded) regions highlight the importance of pre-Columbian food storage techniques (Furquim, 2018; Mendes dos Santos, 2016; Mendes dos Santos et al., 2021), especially for manioc (Fausto & Neves, 2018). One of the products, popularly called Indian bread (*pão de índio* or *pães de índio*), is currently found by chance or in excavations of archaeological sites (Mendes dos Santos et al., 2021). Many techniques have historically been used to produce *pães de índio*. *Pães de índio* are ways of preserving edible plant biomass for a period, especially starch-rich fruits and/or roots. When made with manioc roots, they first processed the manioc to remove anti-nutritional factors (such as prussic acid) and transformed the manioc into a dough, either cooked or raw; fruit mesocarps or seeds were ground into a dough. The dough is shaped, toasted/smoked or not, wrapped in waxy leaves, and buried. Depending upon treatment, *pão de índio* can last for months or years (Mendes dos Santos et al., 2021).

In addition, ceramics are historically used in Amazonia to prepare and store food (Costa et al., 2011). Because ceramics resist degradation in tropical climates, they are more easily found in archaeological sites (Neves, 2011). On the other


hand, leaves, sticks, and vines rot quickly, making them difficult to record in archaeological sites. However, they were undoubtedly widely used, especially in emergencies, such as during extreme floods. Early reports of *pão de índio* considered it a colony of fungi (Araújo & Souza, 1978; Aguiar and Sousa, 1981) or fungus production for food purposes (Santos et al., 2014). However, more recent compositional analyses identified plant materials, mainly edible roots and fruits rich in starch, and their functionality as a form of food storage was confirmed (Mendes dos Santos et al., 2021). Similar to observations in Amazonia, other tropical countries where the manioc is an essential component of local diets use different short, medium, or long-term storage techniques (Knoth, 1993).

There are reports of manioc conservation through burials (Affran, 1968; Anon, 1944; Balagopalan, 2000; Knoth, 1993; Irvine, 1969), covering with various leaves or sawdust (Fadeyibi, 2011; Knoth, 1993; Osei-Opore, 1990; Rickard & Coursey, 1981), and also in plastic bags (Gallat et al., 1998; Knoth, 1993; Rickard & Coursey, 1981; Westby et al., 1999). Manioc storage techniques are believed to be a derivative of the common practice of leaving the roots stored in the field (unharvested) for several months after reaching maturity (Ingram & Humphries, 1972; Rickard & Coursey, 1981). Archeological studies in Amazonia demonstrate that *food storage* techniques were part of the region's ancient local ecological knowledge (LEK) (Furquim, 2018; Neves et al., 2014; Schmidt et al., 2014).

The knowledge about food storage, such as *pães de índio*, is valuable today, keeps changing, and can contribute to the future of local communities (Ávila et al., 2021). Furthermore, it is essential to highlight that changing scenarios, with different social, economic, political, and environmental/ecological pressures, can result in a loss (Reyes-García et al., 2013; Aswani et al., 2018) or changes in LEK (Reyes-García et al. 2014). These pressures influence the dynamics of LEK, a significant area of study in ethnoecology (Gaoue et al., 2017). To manage flood risk, modern adaptation techniques can replace LEK with older techniques (Haughton et al., 2015). In the case of manioc storage, some techniques seem to be in disuse, such as *pão de índio* (Mendes dos Santos, 2021), while modern techniques with the same purpose can take their place. However, different techniques can enhance the resilience of local communities in the context of social, economic, and environmental change (Walker et al., 2004).

Recognizing the great importance of manioc as a source of food and income for *várzea* and *paleovárzea* communities, realizing how extreme flooding can make processing complex, and acknowledging the importance of the dynamics of local knowledge in critical moments, we investigated the techniques known and/or used by local communities

Fig. 1 The study area's map identifies the *várzea* and *paleovárzea* communities in the middle Solimões River region, Amazonas, Brazil



in the *várzea* and *paleovárzea* for storage of manioc roots that allow for further processing and consumption when the water recedes, and whether *várzea* communities know more techniques for storing manioc than *paleovárzea* communities. We hypothesize that: (1) local communities from both ecosystems know and/or use traditional (exclusively made using natural components from local ecosystems, such as fibers, sticks, and leaves, and mentioned in archeological studies about practices and knowledge of Pre-Columbian populations of Amazonia) and modern techniques, that use synthetic components (such as plastic bags) to store the manioc mass, and (2) because *várzea* is more frequently and intensely influenced by flooding, residents in these ecosystems know and use more techniques for storing manioc roots than residents living in the *paleovárzea*.

Methodology

Study Area

This study was conducted in the middle Solimões River basin (Amazonas, Brazil, Fig. 1) between August 2017 and May 2019. Local communities are considered *ribeirinhos*, local populations that historically inhabit the banks of rivers and experience the seasonal variation of waters in their daily lives (CNPCT, 2016).

The *várzeas* originated in the Holocene and are periodically flooded in regular annual cycles, in which soils are covered for up to six months with river waters rich in

sediments with a high content of nutrients from the Andes (Ayres, 2006; Junk et al., 1989). *Paleovárzeas*, on the other hand, originated in the Late Pleistocene and are positioned in the landscape a few meters above the *várzeas*. However, this difference is more significant in regions closer to the mouth of the Amazon River (Irion et al., 2010). Thus, during the annual flood cycle, the *várzea* areas where *ribeirinhos* live and cultivate are usually covered by water for at least 1 to 2 months (Steward et al., 2020). On the other hand, the agricultural and living areas of *paleovárzea* communities are rarely flooded. During extreme flooding, most of the high *várzea* cultivation areas are covered by water, while in the *paleovárzea* only some cultivation areas are affected.

The *várzea* communities in this study are within the Mamirauá Sustainable Development Reserve (RDSM), and the *paleovárzea* communities are within the Amanã Sustainable Development Reserve (RDSA) (Fig. 1). According to the Köppen-Geiger system, the region's climate is tropical rainforest (Af), monthly rainfall is greater than 60 mm throughout the year (Peel et al., 2007), and the average rainfall is 2.200 mm (Ayres, 2006). The flood season (winter) is from December to May, and the dry season (summer) is from June to November (Bueno et al., 2019). On average, the water level of the Solimões River in the study region rises 10.6 m during the flood season and may rise by 15–17 m in more severe floods (Ramalho et al., 2010).

We conducted our study in six communities, three *ribeirinho várzea* communities (Vila Alencar (composed by 13 families), Sítio Fortaleza (14 families) and Porto Braga (33 families) and three *ribeirinho paleovárzea* communities

(*Santa Luzia do Juazinho* (23 families), *Bom Jesus do Calafate* (15 families) and *Santa Luzia do Baré* (6 families)). Information on the socioeconomic characteristics of the residents can be obtained from Ávila et al., (2021). Planting in the region is based on the cultivation of manioc in small areas, combined with the cultivation of other annual and perennial crops (Rognant & Steward, 2015).

Data Collection

We conducted semi-structured interviews with one household head per house until all available families were sampled, totaling 52 *ribeirinhos* in the *várzea* and 36 *ribeirinhos* in the *paleovárzea*. We interviewed one man or woman, more than 18 years old, per family unit. The semi-structured interviews focused on the storage of manioc in years of extreme flooding. For this, we asked if *ribeirinhos* knew any technique for storing manioc. In the case of a positive answer, we asked which techniques they knew. Some specific questions to understand the permanence or continuity of these techniques: (a) whether the *ribeirinho* only knew about the technique but did not know how to perform it, (b) whether the *ribeirinho* knew the technique sufficiently to perform it, and (c) whether the *ribeirinho* had already performed the technique (currently or in the past).

Data Analysis

Information about known and used storage techniques was categorized and analyzed using descriptive statistics. We used a generalized linear model with a binomial distribution using the program R (R Development Core Team, 2020) to verify if there are statistical differences in the *ribeirinhos'* knowledge about manioc storage between the *várzea* and *paleovárzea* ecosystems.

Results

Manioc Storage Techniques

In both the *paleovárzea* and the *várzea*, people from all communities reported that during significant floods they employ various storage techniques to store raw peeled manioc roots (locally called *massa da mandioca*/manioc mass) during the flood and process them into flour when waters recede. They say these techniques were practiced historically and learned from their grandparents and indigenous peoples. The guiding principle of most manioc storage techniques is the creation of an anoxic environment in order to eliminate or reduce oxygen. Four techniques were described in

the study area: burial of manioc mass, basketing of manioc mass, bagging of manioc mass, and *kanaká* of manioc *puba*.

The first technique uses soil and leaves to create a more anoxic environment, called *enterrar a massa da mandioca* (burial of manioc mass). It consists of *ribeirinhos* burying the freshly peeled roots in pits excavated in the soil, which will remain submerged during the flood. The interviewees recognized this technique as widespread and ancient, practiced both in the *várzea* and *paleovárzea*. The technique follows these steps: (1) a 1-meter square pit is dug in the ground (depending on the amount of manioc to store, it can be larger or smaller), (2) the inner walls of the pit are lined with *cauaçú* (*Calathea lutea* (Aubl.) Schult.) leaves, which serve to prevent the roots from direct contact with the soil (one of the respondents mentioned that *embaúba* (*Cecropia* sp.) leaves may also be used for this purpose), (3) peeled manioc roots are placed in the pit up to the edge of the soil surface, (4) *cauaçú* leaves are used to cover all of the manioc roots, and (5) soil excavated from the pit is piled over the manioc to cover the storage pit (Fig. 2).

The second technique, *empaneirar a massa da mandioca* or *empalhar a massa da mandioca* (basketing or wrapping manioc mass), is also locally recognized as ancient and known in both ecosystems; it is cited as a simplified alternative to the first technique. It consists of (1) constructing a fiber basket (popularly called *paneiro*) using dry petioles of *cauaçú* or some *Arecaceae*, (2) lining the fiber basket with *cauaçú* leaves or other leaves, such as *pariri* (cf. *Pouteria pariry* (Ducke) Baehni), (3) filling the basket with peeled manioc roots, (4) covering the top of the basket with *cauaçú* leaves, (5) tying the covering tightly with fiber rope (made using hanging aerial roots of *ambé* (cf. *Philodendron* spp.)), and (6) immersing the basket in the water. When employing this technique, the basket must be completely submerged during the entire storage period (Fig. 2). One of the interviewees also mentioned that the basket could be buried as in the first technique. Others recommended using a rope to support the basket on a tree without touching the river bottom because the manioc stored precisely where the fiber basket touches the river may show rotting signs.

Because major floods often rise quickly, the considerable work required for mass storage of manioc in pits, baskets, or cones may be impractical. About 30 years ago, residents adapted the basketing technique with woven plastic-fiber bags. It is essential that the bag be of woven fiber and not solid plastic since woven fiber bags allow water movement into and out of the bag. This variation is called *ensacamento da massa da mandioca* (bagging of manioc mass). The bagging consists of (1) placing the whole peeled manioc in woven fiber bags, (2) closing the bag with a rope, (3) with another rope, hanging the bag on a tree so that the bag is




Fig. 2 Schematic drawings of techniques used for manioc storage during large floods in *ribeirinho* communities in *várzea* and *paleo-várzea* areas in the middle Solimões River region, Amazonas, Brazil.

completely immersed in the water and does not touch the riverbed during storage (Fig. 2).

A fourth technique, locally called “*kanaká*” (Fig. 2), different from the techniques described above, uses ground manioc root mass (locally called *pubar a mandioca*), which is prepared in a sunken canoe along the river edge where the manioc roots are submerged in water for 2–4 days. As *kanaká* uses only natural components for its construction

(1) Burial of manioc; (2) Manioc basketing; (3) Manioc bagging; (4) Peeled manioc roots wrapped in *cauaçu* leaves for storage in *Kanaká* and *Kanaká* submerged in water (Figures created by Maurício Afonso)

and was cited as used in the past, it was considered an ancient technique to preserve manioc. The *kanaká* steps include (1) in a region that will be flooded, build a cone with *tacana* sticks (cf. Poaceae) fixed to the ground, which is tied with *ambé* vine (cf. *Philodendron* spp.), (2) line the *kanaká* structure with *cauaçu* leaves, (3) place the wet ground manioc mass in the *kanaká*, (4) cover the *kanaká* with *cauaçu* leaves. In this technique, the wet ground manioc mass is

Table 1 Techniques of manioc storage are described in *várzea* and *paleovárzea* communities along the middle Solimões River in Central Amazonia, Brazil

| | Cited in archaeological or ethnographic studies in Amazonia | Material required | Manioc preparation | Labor time | Duration of preservation | Threats like predation | Capacity | Quality of flour |
|---------------------------------|---|--|---|----------------|--------------------------|------------------------|----------|------------------|
| Burial | Yes | <i>Cauaçú</i> leaves | Fresh peeled roots | Intermediate | At least 6 months | Low | High | Low |
| Basket (touching river bottom) | Yes | Fiber basket, fiber rope and <i>cauaçú</i> leaves | Fresh peeled roots | Intermediate | At least 6 months | Intermediate | Low | Low |
| Basket (free in current) | Yes | Fiber basket, fiber rope and <i>cauaçú</i> leaves | Fresh peeled roots | Intermediate | At least 6 months | Intermediate | Low | Intermediate |
| Bagging (touching river bottom) | No | Woven fiber bags | Fresh peeled roots | Fast | At least 6 months | Intermediate | Low | Low |
| Bagging (free in current) | No | Woven fiber bags | Fresh peeled roots | Fast | At least 6 months | Intermediate | Low | Intermediate |
| <i>Kanaká</i> | No | <i>Tacana</i> sticks, <i>ambé</i> and <i>cauaçú</i> leaves | Manioc root mass (manioc “ <i>puba</i> ”) | Time consuming | At least 6 months | Intermediate | High | Intermediate |

isolated from oxygen by the *cauaçú* leaves, allowing storage without oxidation and rot. In a large *kanaká*, manioc can be stored from an entire field (about 0,5–1.0 ha). After processing, such a *kanaká* yields between 8–10 bags of manioc flour- weighing 60 kg each.

The *kanaká* technique was mentioned only by one *ribeirinho* from the *várzea*. He was the oldest man interviewed (63 years) in *Sítio Fortaleza*, with greater traditional knowledge about the community. He was born in a municipality close to the study area (Fonte Boa) and remembered doing these techniques when he was young. However, in other projects conducted by the *Mamirauá* Institute, the *kanaká* practice was observed in other local communities along the middle Solimões River. As our interviews were not conducted in those communities, other mentions about *kanaká* in the region were not included in the results and figures of the present manuscript.

The techniques described above differ in difficulty, speed, and cost (Table 1). Manioc bagging is the fastest, easiest and cheapest technique, as it reuses bags used to pack manioc flour or other plant products. In the case of burial, the leaves must be collected, and there is more physical labor to dig the pit and bury the manioc. The basket, in addition to requiring the collection of leaves, sticks, and ties, also demands the ability to weave the fiber basket and sometimes bury or elevate it with a rope. Finally, the most time-consuming and laborious technique is the *kanaká*. It requires the collection of sticks, ties, and leaves and the structure’s construction.

The manioc mass must also be softened and ground, which takes at least three days to prepare.

Storage time varies with each technique, spanning the time from when cultivation areas are flooded, and the period they need to rebuild the manioc flour houses, which can vary from 1 to 4 months. In the scenario where extreme floods last longer, the *ribeirinhos* confirm that any manioc storage technique mentioned is effective and increasingly necessary.

Ribeirinhos interviewed mentioned that the flour from the stored roots can turn bitter. Because of this, they adopt specific processing techniques to remove the bitter taste. The primary way to do this is to place the previously stored manioc in a sunken canoe near the riverbank, letting the river water soak the roots and wash away the bitterness (in the same way they traditionally soften manioc to prepare the flour). Another strategy to help to remove the bitterness is to let the rainwater wash it off. Even when done effectively, interviewees mention that the flour produced for stored manioc is mainly reserved for family consumption, as its flavor and color can be less acceptable in the market.

Distribution of Knowledge in Different Ecosystems

Among the 88 people interviewed, 32% did not know any technique for storing manioc roots, 20% knew a little about it, 12% knew it in detail but had never practiced it, and 36% knew some technique and had performed one in the past. These proportions, however, varied substantially between

Fig. 3 Percentage of *ribeirinhos* from both ecosystems in the middle Solimões River region, Amazonas, Brazil, who know and/or use manioc storage techniques during extreme floods





Fig. 4 Distribution of knowledge (%) about each manioc storage technique in the study ecosystems in the middle Solimões River region, Amazonas, Brazil, considering only people who know at least one technique



environments; in the *paleovárzea*, most of the interviewees (58%) had performed some technique; in the *várzea* most of the interviewees either did not know any technique (46%) or knew little about it (29%). In this sense, *ribeirinhos* from *paleovárzea* communities use and know more about traditional techniques (72%) than those from *várzea* communities (25%) (Fig. 3).

In general, *ribeirinhos* from the *paleovárzea* know more about traditional manioc storage techniques (87%) than those from the *várzea* (59%) (Fig. 4). In the *várzea* they know mainly the burial technique (47%) and basketing (35%), while in the *paleovárzea* the bagging technique is better known (64%), and the burial technique is used (29%). As for the knowledge, the bagging technique is the best known, followed by the burial technique, the basketing technique, and the *kanaká*.

Among the known techniques, the *kanaká* was mentioned by only one *ribeirinho* in the *várzea*. The burial and basketing techniques are known by at least one *ribeirinho* from each community in both ecosystems. In contrast to what we hypothesized, the results of the generalized linear model to test the knowledge about manioc storage related to the *várzea* and *paleovárzea* ecosystems indicate that in the *paleovárzea* *ribeirinhos* know more about storing manioc than *ribeirinhos* in the *várzea* ($\beta = -1.45$, $SE = 0.53$, $z = -2.76$, $p = 0.005$).

Discussion

Four manioc storage techniques were cited as applicable in the case of extreme and unpredictable flooding. Two of these techniques have not yet been described in the literature for

the region: *kanaká* and *ensacamento* (bagging). One consists of burying manioc roots, and the other three consist of ways in which the peeled manioc is stored in different fiber packages (natural or synthetic), which are submerged in water (basketing, bagging, and *kanaká*). One technique was exclusively mentioned in the *várzea* (*kanaká*), and only the most recently developed technique (bagging) is currently used in both ecosystems. The most significant knowledge dissemination and manioc root storage were in the *paleovárzea*. In climate change, in which extreme flooding is more intense and unpredictable, *ribeirinhos* from both ecosystems, which do not know the techniques were very interested in learning about them. The technique is more used today, given the last extreme flood (2015).

Around the world, local communities also use traditional food storage techniques to preserve potatoes, maize, rice, cowpea, millet, quinoa, sorghum, etc. For example, they use underground pits, wooden structures/silos, or bags (jute, sisal, local grass, cotton, or polyethylene) (Tapia & Torre, 1998; Mobolade et al., 2019).

Manioc storage techniques using burials are cited in old (Edmondson, 1922; Lancaster et al., 1982; Monteiro, 1963) and recent studies in Amazonia (Acosta Muñoz et al., 2005; Mendes dos Santos et al., 2021). Archaeological and ethnographic work highlights the long history of this burial technique in the region who could be used to store starchy plant materials that would be used after the temporary migrations of indigenous groups (Furquim, 2018). Burial is also a technique historically used in Africa (Afran, 1968; Irvine, 1969) and on Mauritius island (Anon, 1944). In the latter case, it is referred to as “Reine’s method,” where layers of peeled manioc roots are alternated with layers of 7.5 cm of earth and then covered with earth (15 cm deep), allowing storage for more than one year (Anon, 1944). In India, there are also studies of manioc burials, where it was observed that after two months in soil with 15% moisture, 80–85% of the roots were undamaged and could be used (Balagopalan, 2000).

As for the basketing technique, use was reported among indigenous groups from other traditional populations of the Solimões River (Sousa et al., 2017) and the upper Negro River (Maia, 2018). In some regions of Hispanic America, where the *cauaçú* also occurs (also known locally as *bijão*), leaves are used to wrap food and protect it from humidity (Sousa et al., 2017). In an archaeological study in the Purus River basin, the authors reported that the basketing technique was used to produce *pães de índio* in the interfluvium of the Jacaréuba-Mucum Rivers due to its specific conservation conditions (Cangussu & Perez, 2017). Concerning the current use of burial and basketing techniques, other studies observed evidence of these practices near the middle Purus River, the upper Negro River, and the upper Madeira River, suggesting that these techniques continue to be used,

although infrequently (Mendes dos Santos et al., 2021). In addition, the historical importance of the burial and basketing techniques has been highlighted for the storage of other roots (such as the *mairá* potato (*Casimirella* sp.)) and also fruits (such as *pupunha* (*Bactris gasipaes* Kunth) and *buriti* (*Mauritia flexuosa* L. f.)) (Lancaster & Coursey, 1984; Mendes dos Santos et al., 2021).

Concerning the *kanaká*, the *puba* process causes the manioc to ferment. This process is also used for manioc flour preparation (locally called *farinha d’água*); the fermented softened roots facilitate the grinding into a homogeneous mass and liberate part of the prussic acid (Lancaster et al., 1982). For the *kanaká* storage technique, some similar methods were observed in Uganda: wooden boxes with damp sawdust. The box is coated with plastic to prevent the sawdust from drying out for 4–8 weeks (Nahdy & Odong, 1995). However, in the case of *kanaká* the wooden structure has a cone shape, the plastic is substituted with local leaves, and manioc *puba* is used.

Storage in conical piles was observed in traditional indigenous silos in Colombia. In these, a thick layer of straw is placed, and the roots are piled up. The mounds, with between 300 and 500 kg of manioc, are covered with straw and earth, with openings left at the bottom for ventilation. The method allows storage for about one month, depending on the temperature and rainfall of the season (Rickard & Coursey, 1981). In the Philippines, conical piles are also used to store manioc for 25 days (Baybay, 1922).

The use of bags to preserve manioc roots is recognized as an extension of traditional storage methods, where plastic bags are used to prevent moisture loss and water stress (Rickard & Coursey, 1981). With bagging, tightly woven bags without chemical treatment were observed in Ghana (Gallat et al., 1998) and Tanzania (Westby et al., 1999), allowing 7–10 days of storage. Polyethylene bags were also used with damp sawdust associated with chemical treatments, preventing post-harvest deterioration and increasing root quality (Carvalho et al., 1985). In Colombia, this type of storage allowed conservation for 14 days (Best, 1990). Comparing the techniques present in the literature with those mentioned in the communities, we can identify that bagging and basketing have similarities with the other techniques of storing manioc in water. Compared to older techniques to store manioc, the bagging technique is more manageable, fast, and practical and demonstrates local knowledge dynamics (Haughton et al., 2015). However, alternatives and old techniques provide options to store manioc, and the knowledge about different techniques are essential factors in enhancing resilience in climate change (Altieri et al., 2015).

Várzeas are at high risk for major periodic flooding (Denevan, 1996). In the *várzea* areas studied, *ribeirinhos*

emphasized that they do not cultivate as much manioc as they used to because extreme floods are recurrent and unpredictable (Avila et al., 2021). Consequently, in *várzea* ecosystems, as there is less manioc to be processed into flour, the need for these storage techniques is lesser compared to the past or the *paleovárzea*. This may be why *várzea ribeirinhos* know less about manioc storage than *ribeirinhos* in the *paleovárzea*.

Conclusion

Ribeirinhos of *várzea* and *paleovárzea* communities are familiar with four techniques for storing fresh manioc. However, in both ecosystems, people used only a recently adapted technique (manioc bagging). *Ribeirinhos* from the *paleovárzea* know more about storage techniques than those from the *várzea*. In the *paleovárzea*, crops do not suffer from annual flooding; thus, the swiddens are more important for income generation and, consequently, are more significant, which could justify that knowledge of this technique is better distributed in this ecosystem. The differences in LEK are due to their histories of interaction with each ecosystem's environmental characteristics, especially susceptibility to extreme flooding. In addition, many *ribeirinhos* in the study area want to know more about manioc storage techniques to avoid losing their crops. This demand illustrates how traditional and adapted technologies are dynamic, need to be shared, and can be crucial in strengthening communities' food security and sovereignty in critical periods, favoring the resilience of these communities even in the context of climate change.

Acknowledgements We thank the residents of the collaborating communities who kindly shared their knowledge with us and strongly supported the study and its logistics. Thanks to the boatmen, field staff, and logistics teams, the Territorialities and Socio-Environmental Governance in the Amazon Research Group, the Archaeology Research Group, and the Agro-ecosystem Management Program, all at the Mamirauá Institute, and the Group for Interdisciplinary Environmental Studies (UFSC-Brazil). CRC thanks the Brazilian Research Council for a research fellowship (PQ 303477/2018-0).

Authors' contributions Conceptualization: Julia Vieira da Cunha Ávila, Charles R. Clement, Angela May Steward; Methodology: Julia Vieira da Cunha Ávila, Angela May Steward; Formal analysis and investigation: Julia Vieira da Cunha Ávila, André Braga Junqueira, Tamara Ticktin; Writing - original draft preparation: Julia Vieira da Cunha Ávila, Charles R. Clement; Writing - review and editing: Anderson Márcio Amaral; André Braga Junqueira; Gilton Mendes dos Santos, Tamara Ticktin; Supervision: Angela May Steward, Charles R. Clement.

Funding CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) provided Julia's PhD Scholarship No 140049/2019-3 and grant agreement No 435985/2018-3, Fundo Amazônia, FAPEAM (Fundação de Amparo à Pesquisa do Estado do Amazonas) under grant

agreement No 062.00148/2020, Finance Code 001 CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior). ABJ has received funding from the European Research Council (ERC) through the LICCI project (ERC Consolidator Grant FP7-771056-LICCI). This work contributes to the "Maria de Maeztu Unit of Excellence" (CEX2019-000940-M).

Data Availability The datasets generated and/or analysed during the current study are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest The authors declare no conflict of interests.

Informed consent The interviews were conducted after the interviewees signed the informed consent form wherein they agreed to participate in the study and authorized the disclosure of the results. The Ethics Committee approved this research for Research with Humans at Mamirauá Institute (CEP authorization number: 2.964.758), registered in the National System for the Management of Genetic Heritage and Traditional Knowledge (SISGEN A494ADE), and obtained authorization from the System of Authorization and Information on Biodiversity (SISBIO 65374-1).

References

- Acosta Muñoz, L. E., Gonzáles, H. A. C., Venegas, C. P. P., Garcia, J. C. A., Arguelles, J., Valderrama, A. M., & Huaines, F. J. (2005). La práctica de conservación de biomasa de yuca: tecnología tradicional indígena aplicada en el presente. In L. E. Acosta Muñoz & A. M. Valderrama (Eds.), *Enterramientos de masas de yuca del pueblo Ticuna: Tecnología tradicional en la várzea del Amazonas colombiano* (pp. 63–73). Instituto Amazónico de investigaciones Científicas, Bogotá: #225
- Adams, C., Murrieta, R., Siqueira, A., Neves, W., & Sanches, R. (2009). Bread of the land: The invisibility of manioc in the Amazon. In C. Adams, R. Murrieta, W. Neves, & M. Harris (Eds.), *Amazon peasant societies in a changing environment* (pp. 281–305). Springer. https://doi.org/10.1007/978-1-4020-9283-1_13. Dordrechts
- Affran, D. (1968). Cassava and its economic importance. *Ghana Farmer*, 12(4), 172–178
- Aguiar, I. de J. A., & Sousa, M. A. de. (1981). Polyporus indigenus I. Araujo & M. A. Sousa, nova espécie da Amazônia. *Acta Amazonica*, 11(3), 449–455. <https://doi.org/10.1590/1809-43921981113449>
- Alencar, E. F. (2010). *Memórias de Mamirauá*. Instituto de Desenvolvimento Sustentável Mamirauá
- Altieri, M. A., Nicholls, C. I., Henao, A., & Lana, M. A. (2015). Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*, 35(3), 869–890. <https://doi.org/10.1007/s13593-015-0285-2>
- Anon. (1944). La conservation du manioc par le procede de Reine. *Revue Agricole et Sucrière de l'île Maurice*, 23(3), 105–106
- Araújo, I. J., & Souza, M. A. (1978). Nota prévia sobre o pão do índio da Amazônia Brasileira. *Acta Amazonica*, 8(2), 316–318
- Aswani, S., Lemahieu, A., & Sauer, W. H. H. (2018). Global trends of local ecological knowledge and future implications. *PLOS ONE*, 13(4), e0195440. <https://doi.org/10.1371/journal.pone.0195440>
- Ávila, J. V. C., Clement, C. R., Junqueira, A. B., Ticktin, T., & Steward, A. M. (2021). Adaptive management strategies of local communities in two Amazonian floodplain ecosystems in the face of

- extreme climate events. *Journal of Ethnobiology*, 41(3), 409–426. <https://doi.org/10.2993/0278-0771-41.3.409>
- Ayres, J. M. (2006). *As matas de várzea do Mamirauá: Médio Rio Solimões* (3rd ed.). Tefé: Sociedade Civil Mamirauá
- Balagopalan, C. (2000). *Integrated technologie for value addition and post harvest management in tropical tuber Crops*. Central Tuber Crops Research Intitute
- Baybay, D. S. (1922). Storage of some root crops and other perishable farm products. *Philippine Agriculture*, 10(9), 423–440
- Best, R. (1990). Storage and processing of cassava in Latin America, lessons for the subregion (West Africa). *Action programme for the prevention of food losses, improving post-harvest handling, storage and processing of root and tuber Crops* (pp. 9–20). FAO
- Bueno, G. T., Cherem, L. F. S., Toni, F., Guimarães, F. S., & Bayer, M. (2019). *Amazonia* (pp. 169–197). https://doi.org/10.1007/978-3-030-04333-9_9
- Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M. J., Wu, L., England, M. H., Wang, G., Guilyardi, E., & Jin, F. F. (2014). Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature Climate Change*, 4(2), 111–116. <https://doi.org/10.1038/nclimate2100>
- Cangussu, D., & Perez, W. (2017). *Relatório da expedição de localização Jacareituba*. FUNAI, CGIIRC, FPE Madeira-Purus
- de Carvalho, V. D., Chalfoun, S. M., & Juste Junior, E. S. G. (1985). Storage methods on fresh cassava roots preservation, 1: effect of polyethylene bags and humid sawdust associated to chemical treatments on post-harvest deterioration and root quality. *Revista Brasileira de Mandioca (Brazil)*, 4(1), 79–85
- CNPCT (2016). Comissão Nacional de Desenvolvimento Sustentável de Povos e Comunidades Tradicionais. <http://Portalyfade.Mma.Gov.Br/Ribeirinhos-Caracteristicas/120-Povos-e-Comunidades/Ribeirinhos>.
- Costa, M. L., da Rios, G. M., Silva, M. M. C., da Silva, G. J. da, & Molano-Valdes, U. (2011). Mineralogy and chemistry of archaeological ceramic fragments from archaeological Dark Earth site in Colombian Amazon. *Revista Escola de Minas*, 64(1), 17–23. <https://doi.org/10.1590/S0370-44672011000100002>
- Denevan, W. M. (1996). A bluff model of riverine settlement in pre historic Amazonia. *Annals of the Association of American Geographers*, 86(4), 654–681
- Dubreuil, V., Funatsu, B. M., Michot, V., Nasuti, S., Debortoli, N., de Mello-Thery, N. A., & Le Tourneau, F. M. (2017). Local rainfall trends and their perceptions by Amazonian communities. *Climatic Change*, 143(3–4), 461–472. <https://doi.org/10.1007/s10584-017-2006-0>
- Dufour, D. L. (1995). A closer look at the nutritional implications of bitter cassava use. In: Sponsel LE, ed. *Indigenous peoples and the future of Amazônia*. In L. Sponse (Ed.), *Indigenous peoples and the future of Amazônia*. (pp. 149–165). University of Arizona Press
- Dufour, D. L., Piperata, B. A., Murrieta, R. S. S., Wilson, W. M., & Williams, D. D. (2016). Amazonian foods and implications for human biology. *Annals of Human Biology*, 43(4), 330–348. <https://doi.org/10.1080/03014460.2016.1196245>
- Edmondson, G. (1922). *Journal of the Travels and Labours of Father Samuel Fritz in the River of the Amazons between 1686 and 1723*. Haklayt Soc
- Fadeyibi, A. (2011). Storage methods and some uses of cassava in Nigeria. *Continental Journal of Agricultural Science*, 5(2), 12–18
- Fausto, C., & Neves, E. G. (2018). Was there ever a Neolithic in the Neotropics? Plant familiarization and biodiversity in the Amazon. *Antiquity*, 92(366), 1604–1618. <https://doi.org/10.15184/ajcc.2018.157>
- Fraser, J. A., Alves-Pereira, A., Junqueira, A. B., Peroni, N., & Clement, C. R. (2012). Convergent adaptations: bitter manioc cultivation systems in fertile anthropogenic dark earths and floodplain soils in Central Amazonia. *Plos One*, 7(8), e43636. <https://doi.org/10.1371/journal.pone.0043636>
- Funatsu, B. M., Dubreuil, V., Racapé, A., Debortoli, N. S., Nasuti, S., & Le Tourneau, F. M. (2019). Perceptions of climate and climate change by Amazonian communities. *Global Environmental Change*, 57. <https://doi.org/10.1016/j.gloenvcha.2019.05.007>
- Furquim, L. (2018). *Arqueobotânica e mudanças socioeconômicas no Sudoeste Amazônico durante o Holoceno Médio*. Universidade de São Paulo
- Gallat, S., Crentsil, D., & Bancroft, R. D. (1998). Development of a low cost cassava fresh root storage technology for the Ghanaian market. In R. S. B. Ferris (Ed.), *Postharvest technology and commodity marketing. Proceedings of a post-harvest conference*. Accra, Ghana. (pp. 77–84)
- Gaoue, O. G., Coe, M. A., Bond, M., Hart, G., Seyler, B. C., & McMillen, H. (2017). theories and major hypotheses in ethnobotany. *Economic Botany*, 71(3), 269–287. <https://doi.org/10.1007/s12231-017-9389-8>. Springer New York LLC
- Haughton, G., Bankoff, G., & Coulthard, J., T (2015). In search of ‘lost’ knowledge and outsourced expertise in flood risk management. *Transactions of the Institute of British Geographers*, 40(3), 375–386. <https://doi.org/10.1111/tran.12082>
- Ingram, J. S., & Humphries, J. R. O. (1972). Cassava storage - a review. *Tropical Science*, 14, 131–148
- Irion, G., de Mello, J. A. S. N., Morais, J., Piedade, M. T. F., Junk, W. J., & Garming, L. (2010). Development of the Amazon Valley during the middle to late Quaternary: *Sedimentological and Climatological Observations* (pp. 27–42). https://doi.org/10.1007/978-90-481-8725-6_2
- Irvine, F. R. (1969). *African Crops*. (3rd ea.). Oxford Univ. Press
- Junk, W. J., Bayley, P. B., & Sparks, R. E. (1989). The flood pulse concept in river-floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences*, 106(1), 110–127
- Knoth, J. (1993). *Traditional storage of yams and cassava and its improvement*. GTZ: Postharvest Project
- Lancaster, P. A., & Coursey, D. G. (1984). Traditional post-harvest technology of perishable tropical staples. In *FAO Agricultural Services Bulletin n°59* (p. 88). <http://www.fao.org/3/x5045e/x5045E00.htm#Contents>
- Lancaster, P. A., Ingram, J. S., Lim, M. Y., & Coursey, D. G. (1982). Traditional cassava-based foods: survey of processing techniques. *Economic Botany*, 36(1), 12–45. www.jstor.org/stable/4254349
- Langill, J. C., & Abizaid, C. (2020). What is a bad flood? Local perspectives of extreme floods in the Peruvian Amazon. *Ambio*, 49(8), 1423–1436. <https://doi.org/10.1007/s13280-019-01278-8>
- Lentz, D. L. (2000). Anthropocentric food webs in the Precolumbian Americas. In *Imperfect balance: landscape transformations in the Precolumbian Americas* (pp. 89–119)
- Lima, D., Steward, A., & Richers, B. T. (2012). Trocas, experimentações e preferências: um estudo sobre a dinâmica da diversidade da mandioca no médio Solimões, Amazonas. *Boletim Do Museu Paraense Emílio Goeldi Ciências Humanas*, 7(2), 371–396. <https://doi.org/10.1590/S1981-81222012000200005>
- Maia, G. (2018). *Bahsamori: o tempo, as estações e as etiquetas sociais dos Yepamahsã (Tukano)*. Edua/Neai
- Marengo, J. A., Borma, L. S., Rodriguez, D. A., Pinho, P., Soares, W. R., & Alves, L. M. (2013). Recent extremes of drought and flooding in Amazonia: vulnerabilities and human adaptation. *American Journal of Climate Change*, 02(02), 87–96. <https://doi.org/10.4236/ajcc.2013.22009>
- dos Mendes, G. (2016). Plantas e parentelas: notas sobre a história da agricultura no Médio Purus. In G. Mendes dos Santos, & M. Aparicio (Eds.), *Redes Arawa—Ensaio de etnologia do Médio Purus* (pp. 19–39). Braz.: EDUA

- dos Mendes, G., Cangussu, D., Furquim, L. P., Watling, J., & Neves, E. G. (2021). Pão-de-índio e massas vegetais: elos entre passado e presente na Amazônia indígena. *Boletim Do Museu Paraense Emílio Goeldi Ciências Humanas*, 16(1), <https://doi.org/10.1590/2178-2547-bgoeldi-2020-0012>
- Mobolade, A. J., Bunindro, N., Sahoo, D., & Rajashekar, Y. (2019). Traditional methods of food grains preservation and storage in Nigeria and India. *Annals of Agricultural Sciences*, 64(2), 196–205. <https://doi.org/10.1016/j.aosas.2019.12.003>
- Monteiro, M. Y. (1963). Alimentos preparados à base da mandioca. *Revista Brasileira de Folclore*, III(5), 37–82
- Nahdy, S. M., & Odong, M. (1995). Storage of fresh cassava tuber in plant based storage media. In Accra (Ed.), *Proceeding of the workshop on "post-harvest technology experience in Africa"* (pp. 4–8). FAO
- Neves, E. G. (2011). Archaeological cultures and past identities in the pre-colonial Central Amazon. *Ethnicity in ancient Amazonian: reconstructing past identities from archaeology, linguistic and ethnohistory* (p. 1:27). University Press of Colorado
- Neves, E. G., Guapindaia, V. L. C., Lima, H. P., Costa, B. L. S., & Gomes, J. (2014). A Tradição Pocó-Açutuba e os primeiros sinais visíveis de modificações de paisagens na calha do Amazonas. In S. Rostain (Ed.), *Amazonia, Memórias de las conferencias magistrales del 3er Encuentro Internacional de Arqueologia Amazónica*.
- Osei-Opare, A. F. (1990). Household and village level storage and processing of root and tuber crops in Ghana. Lessons for the Sub-Region West Africa. *Action programme for the prevention of food losses. improving post-harvest handling, storage and processing of root and tuber crops* (pp. 46–57). Food And Agriculture Organization (FAO)
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and earth system sciences*, 11(5), 1633–1644. <https://doi.org/10.5194/hess-11-1633-2007>
- Penteado, I. M., Nascimento, A. C. S., Corrêa, D., Moura, E. A. F., Zilles, R., Gomes, M. C. R. L., Pires, F. J., Brito, O. S., da Silva, J. F., Reis, A. V., Souza, A., & Pacifico, A. C. N. (2019). Among people and artifacts: Actor-Network Theory and the adoption of solar ice machines in the Brazilian Amazon. *Energy research and social science*, 53(February), 1–9. <https://doi.org/10.1016/j.erss.2019.02.013>
- Posey, D. A. (1985). Indigenous management of tropical forest ecosystems: the case of the Kayapó Indians of the Brazilian Amazon. *Agroforestry Systems*, 3(2), 139–158. <https://doi.org/10.1007/BF00122640>
- R Development Core Team (2020). *R: A Language and Environment for Statistical Computing*.
- Ramalho, E. E., Macedo, J., Vieira, T. M., Valsecchi, J., Calvimontes, J., Marmontel, M., & Queiroz, H. L. (2010). Ciclo hidrológico nos ambientes de várzea da Reserva de Desenvolvimento Sustentável Mamirauá – Médio Rio Solimões, período de 1990 a 2008. *UAKARI*, 5(1), 61–87
- Reyes-García, V., Aceituno-Mata, L., Calvet-Mir, L., Garnatje, T., Gómez-Baggethun, E., Lastra, J. J., Ontillera, R., Parada, M., Rigat, M., Vallès, J., Vila, S., & Pardo-de-Santayana, M. (2014). Resilience of traditional knowledge systems: The case of agricultural knowledge in home gardens of the Iberian Peninsula. *Global Environmental Change*, 24, 223–231. <https://doi.org/10.1016/j.gloenvcha.2013.11.022>
- Reyes-García, V., Guèze, M., Luz, A. C., Paneque-Gálvez, J., Macía, M. J., Orta-Martínez, M., Pino, J., & Rubio-Campillo, X. (2013). Evidence of traditional knowledge loss among a contemporary indigenous society. *Evolution and Human Behavior*, 34(4), 249–257. <https://doi.org/10.1016/j.evolhumbehav.2013.03.002>
- Rickard, J. E., & Coursey, D. G. (1981). Cassava storage, part I: storage of fresh cassava roots. *Tropical Science*, 23, 1–32
- Rognant, C., & Steward, A. (2015). Qui garde le mieux la terre?. L'agriculture familiale face aux stratégies de conservation forestière dans la Réserve de Développement Durable Amanã, Amazonas, Brésil. *Anthropology of Food*, S11
- Santos, C., Silva, F. E. C., da Silva, S. F. S. M., da Sullasi, H. L., de Paula, A. S., Oliveira, C. A., Ghett, N. C., & de Castro, V. M. C. (2014). O pão do índio dos Mukini da Amazônia: estudo de caso sobre o uso de suprimentos específicos na dieta alimentar indígena. *Fundação Museu Do Homem Americano*, 1(11), 58–77
- Schmidt, M. J., Py-Daniel, R., de Paula Moraes, A., Valle, C., Caromano, R. B. M., Texeira, C. F., Barbosa, W. G., Fonseca, C. A., Magalhães, J. A., Guapindaia, M. P., Moraes, V. L., Lima, B., Neves, H. P., E. G., & Heckenberger, M. J. (2014). Dark earths and the human built landscape in Amazonia: a widespread pattern of anthrosol formation. *Journal of Archaeological Science*, 42, 152–165. <https://doi.org/10.1016/j.jas.2013.11.002>
- Sousa, M. D. J. S., Bezerra, N. P., Leoni, J. M., Oliveira, M. M. D. C., & Amaral, M. R. A. (2017). Teçume d'Amazônia: fortalecimento político das mulheres produzindo vitalidade de conhecimentos tradicionais. *Amazônica - Revista de Antropologia*, 8(2), 310. <https://doi.org/10.18542/amazonica.v8i2.5046>
- Steward, A. M., Costa-Santos, R. B., Rognant, C., Viana, F. M. F., Ávila, J. V. C., Santos, J. G., Rodrigues, J., & Vieira, S. (2022). Rising Above the Flood: Modifications in Agricultural Practices and Livelihood Systems in Central Amazonia – Perspectives from Ribeirinho and Indigenous Communities. M. Roué, D. Nakashima, & I. Krupnik (Eds.), *Resilience through Knowledge Co-Production: Indigenous Knowledge, Science, and Global Environmental Change* (pp. 233–246). Cambridge: Cambridge University Press. <https://doi.org/10.1017/9781108974349.018>
- Tapia, M. E., & Torre, A. D. (1998). *la. Women farmers and Andean seeds*
- Tregidgo, D., Barlow, J., Pompeu, P. S., & Parry, L. (2020). Tough fishing and severe seasonal food insecurity in Amazonian flooded forests. *People and Nature*, 2(2), 468–482. <https://doi.org/10.1002/pan3.10086>
- Valer, L. R., Mocelin, A., Zilles, R., Moura, E., & Nascimento, A. C. S. (2014). Assessment of socioeconomic impacts of access to electricity in Brazilian Amazon: Case study in two communities in Mamirauá Reserve. *Energy for sustainable development*, 20(1), 58–65. <https://doi.org/10.1016/j.esd.2014.03.002>
- Venturato, R. D., & Pereira, K. J. C. (2010). Aspects of food sovereignty and labor sharing in domestic units at the Mamirauá and Amanã Sustainable Development Reserves. *Scientific Magazine UAKARI*, 6(2), 21–33. <https://doi.org/10.31420/uakari.v6i2.90>
- Walker, B., Holling, C. S., Carpenter, S. R., & Kinzig, A. P. (2004). Resilience, Adaptability and Transformability in Social-ecological Systems. *Ecology and Society*, 9(2), art5. <https://doi.org/10.5751/ES-00650-090205>
- Westby, A., Kleih, U., Hall, A., Ndunguru, G., Crentsil, D., Bockett, G., & Graffham, A. (1999). Needs assessment in post-harvest research and development. In I. F. Grant & C. R. Sear (Eds.), *Decision tools in sustainable development* (pp. 143–160)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.